

Product Datasheet

Order: order@ebiocell.com

TEL: (540)808-3925

Supprt: tech@ebiocell.com
Web: www.ebiocell.com

PARP-1 Rabbit Polyclonal Antibody

Catalog #: EAB10030

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IP, IHC-P, IF/ICC, ChIP	113	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IP(Immunoprecipitation)	1:20-200
IHC-P(Immunohistochemistry-Paraffin)	1:50-300
IF/ICC(Immunofluorescence/Immunocytochemistry)	1:50-300
ChIP(Chromatin Immunoprecipitation)	1:10-100

Product Information

Conjugate Unconjugate

Specificity PARP-1 Rabbit Polyclonal Antibody detects endogenous levels of PARP-1 protein.

Purification Affinity purification

Concentration1mg/mlFormatLiquid

Formulation In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol

Shipping Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks

 UniProt ID
 P09874

 Entrez-Gene Id
 142

Product Description

PARP-1(Poly(ADP-ribose) polymerase-1), also designated PARP, is a nuclear DNA-binding zinc finger protein that influences DNA repair, DNA replication, modulation of chromatin structure, and apoptosis. In response to genotoxic stress, PARP-1 catalyzes the transfer of ADP-ribose units from NAD(+) to a number of acceptor molecules including chromatin. PARP-1 recognizes DNA strand interruptions and can complex with RNA and negatively regulate transcription. Actinomycin D- and etoposide-dependent induction of caspases mediates cleavage of PARP-1 into a p89 fragment that traverses into the cytoplasm. Apoptosis-inducing factor (AIF) translocation from the mitochondria to the nucleus is PARP-1-dependent and is necessary for PARP-1-dependent cell death. PARP-1 deficiencies lead to chromosomal instability due to higher frequencies of chromosome fusions and aneuploidy, suggesting that poly(ADP-ribosyl)ation contributes to the efficient maintenance of genome integrity.