

nNOS/NOS1 Rabbit Polyclonal Antibody

Catalog #: EAB10270

| Host/Isotype | Clonality | Applications | MW (kDa) | Reactivity |
|--------------|------------|--------------------------|----------|-------------------|
| Rabbit IgG | Polyclonal | WB, IHC-P, IF/ICC, ELISA | 160 | Human, Mouse, Rat |

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| | |
|--|--------------|
| WB (Western Blotting) | 1:500-2000 |
| IHC-P (Immunohistochemistry-Paraffin) | 1:50-300 |
| IF/ICC (Immunofluorescence/Immunocytochemistry) | 1:50-300 |
| ELISA (Enzyme-linked Immunosorbent Assay) | 1:5000-20000 |

Product Information

| | |
|-----------------------|---|
| Conjugate | Unconjugate |
| Specificity | nNOS/NOS1 Rabbit Polyclonal Antibody detects endogenous levels of nNOS/NOS1 protein. |
| Purification | Affinity purification |
| Concentration | 1mg/ml |
| Format | Liquid |
| Formulation | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol |
| Shipping | Gel Pack |
| Storage | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks |
| UniProt ID | P29475 |
| Entrez-Gene Id | 4842 |

Product Description

Nitric oxide (NO) has a broad range of biological activities and has been implicated in signaling pathways in phylogenetically diverse species. Nitric oxide synthases (NOSs), the enzymes responsible for synthesis of NO, contain an N-terminal oxygenase domain and a C-terminal reductase domain. NOS activity requires homodimerization as well as three cosubstrates (L-arginine, NADPH and O₂) and five cofactors or prosthetic groups (FAD, FMN, calmodulin, tetrahydrobiopterin and heme). Several distinct NOS isoforms have been described and been shown to represent the products of three distinct genes. These include two constitutive Ca²⁺/CaM-dependent forms of NOS, including NOS1 (also designated ncNOS) whose activity was first identified in neurons, and NOS3 (also designated ecNOS), first identified in endothelial cells. The inducible form of NOS, NOS2 (also designated iNOS), is Ca²⁺-independent and is expressed in a broad range of cell types.

For Reserch Use Only. Not For Use In Diagnostic Procedures