

## ICAM-1/CD54 Rabbit Polyclonal Antibody

### Catalog #: EAB10307

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF, ELISA	58	Human, Mouse, Rat

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IHC-P(Immunohistochemistry-Paraffin)	1:50-300
IF(Immunofluorescence)	1:50-300
ELISA(Enzyme-linked Immunosorbent Assay)	1:5000-20000

### Product Information

Conjugate	Unconjugate
Specificity	ICAM-1/CD54 Rabbit Polyclonal Antibody detects endogenous levels of ICAM-1/CD54 protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	<a href="#">P05362</a>
Entrez-Gene Id	<a href="#">3383</a>

### Product Description

Cell adhesion molecules (CAMs) are a family of closely related cell surface glycoproteins involved in cell-cell interactions during growth and are thought to play important, yet separate, roles in embryogenesis and development. The intracellular adhesion molecule-1 (ICAM-1), also referred to as CD54, is an integral membrane protein of the immunoglobulin superfamily and recognizes the  $\beta 2\alpha 1$  and  $\beta 2\alpha M$  Integrins. This gene encodes a cell surface glycoprotein which is typically expressed on endothelial cells and cells of the immune system. It binds to integrins of type CD11a / CD18, or CD11b / CD18 and is also exploited by Rhinovirus as a receptor.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**