

PDGF Receptor β Rabbit Polyclonal Antibody

Catalog #: EAB10418

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF/ICC, ELISA	124	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IHC-P(Immunohistochemistry-Paraffin)	1:50-300
IF/ICC(Immunofluorescence/Immunocytochemistry)	1:50-300
ELISA(Enzyme-linked Immunosorbent Assay)	1:5000-20000

Product Information

Conjugate	Unconjugate
Specificity	PDGF Receptor β Rabbit Polyclonal Antibody detects endogenous levels of PDGF Receptor β protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	P09619
Entrez-Gene Id	5159

Product Description

The protein encoded by this gene is a cell surface tyrosine kinase receptor for members of the platelet-derived growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer (PDGFB or PDGFD) or a heterodimer (PDGFA and PDGFB). This gene is essential for normal development of the cardiovascular system and aids in rearrangement of the actin cytoskeleton. This gene is flanked on chromosome 5 by the genes for granulocyte-macrophage colony-stimulating factor and macrophage-colony stimulating factor receptor; all three genes may be implicated in the 5-q syndrome. A translocation between chromosomes 5 and 12, that fuses this gene to that of the ETV6 gene, results in chronic myeloproliferative disorder with eosinophilia.

For Reserch Use Only. Not For Use In Diagnostic Procedures