

HSP90 α Rabbit Polyclonal Antibody

Catalog #: EAB10478

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF	90	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:1000-5000
IHC-P(Immunohistochemistry-Paraffin)	1:100-500
IF(Immunofluorescence)	1:50-300

Product Information

Conjugate	Unconjugate
Specificity	HSP90 α Rabbit Polyclonal Antibody detects endogenous levels of HSP90 α protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	P07900
Entrez-Gene Id	3320

Product Description

The heat shock response was first described for *Drosophila* salivary gland cells and morphologically consists of a change in their polytene chromosome puffing patterns that involves de novo synthesis of a few proteins. Similar heat shock proteins were later discovered in bacterial chicken and mammalian cells, and have been subsequently studied in other organisms. A series of proteins, including HSP 90, HSP 70, HSP 20-30 and ubiquitin, are induced by insults such as temperature shock, chemicals and other environmental stress. A major function of HSP 90 and other HSPs is to act as molecular chaperones. HSP 90 forms a complex with glucocorticoid receptor (GR), rendering the non ligand-bound receptor transcriptionally inactive. HSP 90 binds the GR as a heterocomplex composed of either HSP 56 or Cyclophilin D, forming an aporeceptor complex. HSP 90 also exists as a dimer with other proteins such as p60/ST11 and p23, forming an aporeceptor complex with estrogen and androgen receptors.

For Reserch Use Only. Not For Use In Diagnostic Procedures