

Atg4D Rabbit Polyclonal Antibody

Catalog #: EAB10494

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC, IF, ELISA	53	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

IHC-P(Immunohistochemistry-Paraffin)1:50-300IF(Immunofluorescence)1:50-300ELISA(Enzyme-linked Immunosorbent Assay)1:5000-20000	WB(Western Blotting)	1:500-2000
	IHC-P(Immunohistochemistry-Paraffin)	1:50-300
ELISA(Enzyme-linked Immunosorbent Assay) 1:5000-20000	IF(Immunofluorescence)	1:50-300
	ELISA(Enzyme-linked Immunosorbent Assay)	1:5000-20000

Product Information

Conjugate	Unconjugate
Specificity	Atg4D Rabbit Polyclonal Antibody detects endogenous levels of Atg4D protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	<u>Q86TL0</u>
Entrez-Gene Id	<u>84971</u>

Product Description

Autophagy is the process by which endogenous proteins and damaged organelles are destroyed intracellularly. Autophagy is postulated to be essential for cell homeostasis and cell remodeling during differentiation, metamorphosis, non-apoptotic cell death, and aging. Reduced levels of autophagy have been described in some malignant tumors, and a role for autophagy in controlling the unregulated cell growth linked to cancer has been proposed. This gene encodes a member of the autophagin protein family. The encoded protein is also designated as a member of the C-54 family of cysteine proteases.

For Reserch Use Only. Not For Use In Diagnostic Procedures

EbioCell Lifescineces, Inc.

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA