

## TGF-β Receptor II Rabbit Polyclonal Antibody

# Catalog #: EAB10550

| Host/Isotype | Clonality  | Applications             | MW (kDa) | Reactivity        |
|--------------|------------|--------------------------|----------|-------------------|
| Rabbit IgG   | Polyclonal | WB, IHC-P, IF/ICC, ELISA | 65       | Human, Mouse, Rat |

#### **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| WB(Western Blotting)                           | 1:500-2000   |
|--|--------------|
| IHC-P(Immunohistochemistry-Paraffin)           | 1:50-300     |
| IF/ICC(Immunofluorescence/Immunocytochemistry) | 1:50-300     |
| ELISA(Enzyme-linked Immunosorbent Assay)       | 1:5000-20000 |
|  |              |

#### **Product Information**

| Conjugate      | Unconjugate   |
|----------------|---|
| Specificity    | TGF-β Receptor II Rabbit Polyclonal Antibody detects endogenous levels of TGF-β Receptor II<br>protein.                                   |
| Purification   | Affinity purification   |
| Concentration  | 1mg/ml  |
| Format         | Liquid  |
| Formulation    | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol  |
| Shipping       | Gel Pack  |
| Storage        | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles.<br>Aliquots may be stored at +4°C for 1-2 weeks |
| UniProt ID     | <u>P37173</u>   |
| Entrez-Gene Id | 7048  |

### **Product Description**

The protein encoded by this gene is a transmembrane protein that has a protein kinase domain, forms a heterodimeric complex with TGFbeta receptor type-1, and binds TGF-beta. This receptor/ligand complex phosphorylates proteins, which then enter the nucleus and regulate the transcription of genes related to cell proliferation, cell cycle arrest, wound healing, immunosuppression, and tumorigenesis. Mutations in this gene have been associated with Marfan Syndrome, Loeys-Deitz Aortic Aneurysm Syndrome, and the development of various types of tumors. Alternatively spliced transcript variants encoding different isoforms have been characterized.

For Reserch Use Only. Not For Use In Diagnostic Procedures

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA