

Product Datasheet

Order: order@ebiocell.com

TEL: (540)808-3925 tech@ebiocell.com

Supprt: tech@ebiocell.com
Web: www.ebiocell.com

Ubiquitin/Ub Mouse Monoclonal Antibody

Catalog #: EAB10565

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Mouse IgG1	Monoclonal	WB, IHC-P, IF/ICC		Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting) 1:500-2000
IHC-P(Immunohistochemistry-Paraffin) 1:50-300
IF/ICC(Immunofluorescence/Immunocytochemistry) 1:50-300

Product Information

Conjugate Unconjugate

Specificity Ubiquitin/Ub Mouse Monoclonal Antibody detects endogenous levels of Ubiquitin/Ub protein.

Purification Affinity purification

Concentration1mg/mlFormatLiquid

Formulation In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol

Shipping Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks

UniProt ID <u>P62987</u>, <u>P0CG48</u>, <u>P0CG47</u>, <u>P62979</u>

Entrez-Gene Id <u>7311, 7316, 7314, 6233</u>

Product Description

This gene encodes ubiquitin, one of the most conserved proteins known. Ubiquitin has a major role in targeting cellular proteins for degradation by the 26S proteosome. It is also involved in the maintenance of chromatin structure, the regulation of gene expression, and the stress response. Ubiquitin is synthesized as a precursor protein consisting of either polyubiquitin chains or a single ubiquitin moiety fused to an unrelated protein. This gene consists of three direct repeats of the ubiquitin coding sequence with no spacer sequence. Consequently, the protein is expressed as a polyubiquitin precursor with a final amino acid after the last repeat. An aberrant form of this protein has been detected in patients with Alzheimer's disease and Down syndrome. Pseudogenes of this gene are located on chromosomes 1, 2, 13, and 17. Alternative splicing results in multiple transcript variants.