

Phospho-TACC3 (Ser558) Rabbit Polyclonal Antibody

Catalog #: EAB10597

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB	90	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting) 1:500-2000

Product Information

Conjugate	Unconjugate
Specificity	Phospho-TACC3 (Ser558) Rabbit Polyclonal Antibody detects endogenous levels of TACC3 protein only when phosphorylated at Ser558.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	Q9Y6A5
Entrez-Gene Id	10460

Product Description

TACC3 also designated ERIC-1 is a motor spindle protein that may play a role in stabilization of the mitotic spindle. This protein may also play a role in growth a differentiation of certain cancer cells. TACC family members, TACC1, TACC2 and TACC3, map very closely to the corresponding FGFR1, FGFR2 and FGFR3 genes on chromosomes 8, 10 and 4. Subsequently, since they are phylogenetically related, it is proposed that TACC and FGFR have similar roles in cell growth and differentiation. Also, TACC1 contains a conserved C-terminal region as in the Drosophila homolog, D-TACC. It has been shown that D-TACC is necessary for normal spindle function, and the mammalian TACC proteins appears to interact with centrosomes and microtubules in a similar manner.

For Reserch Use Only. Not For Use In Diagnostic Procedures