

**Product Datasheet** 

Order: order@ebiocell.com

TEL: (540)808-3925

Supprt: tech@ebiocell.com
Web: www.ebiocell.com

## A-Raf Rabbit Polyclonal Antibody

Catalog #: EAB10624

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF, ELISA	68	Human, Mouse, Rat

## **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)1:500-2000IHC-P(Immunohistochemistry-Paraffin)1:50-300IF(Immunofluorescence)1:50-300ELISA(Enzyme-linked Immunosorbent Assay)1:5000-20000

## Product Information

**Conjugate** Unconjugate

**Specificity** A-Raf Rabbit Polyclonal Antibody detects endogenous levels of A-Raf protein.

**Purification** Affinity purification

Concentration1mg/mlFormatLiquid

Formulation In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol

Shipping Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks

UniProt ID P10398
Entrez-Gene Id 369

## **Product Description**

Several serine/threonine protein kinases have been implicated as intermediates in signal transduction pathways. These include ERK/MAP kinases, ribosomal S6 kinase (Rsk) and Raf-1. Raf-1 is a cytoplasmic protein with intrinsic serine/threonine activity. It is broadly expressed in nearly all cell lines tested to date and is the cellular homolog of v-Raf, the product of the transforming gene of the 3611 strain of murine sarcoma virus. The unregulated kinase activity of the v-Raf protein has been associated with transformation and mitogenesis while the activity of Raf-1 is normally suppressed by a regulatory N-terminal domain. A-Raf, a second member of the Raf gene family of serine/threonine protein kinases, exhibits substantial homology to Raf-1 within the kinase domain of the two molecules, but less homology elsewhere. Expression of A-Raf is found at highest levels in urogenital tissues and kidney and at lowest level in brain tissue.