

Phospho-Catenin δ -1 (Tyr228) Rabbit Polyclonal Antibody

Catalog #: EAB10669

| Host/Isotype | Clonality | Applications | MW (kDa) | Reactivity |
|--------------|------------|------------------------|----------|-------------------|
| Rabbit IgG | Polyclonal | WB, IHC, IF/ICC, ELISA | 108 | Human, Mouse, Rat |

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| | |
|--|---------------|
| WB (Western Blotting) | 1:500-2000 |
| IHC (Immunohistochemistry) | 1:50-300 |
| IF/ICC (Immunofluorescence/Immunocytochemistry) | 1:50-300 |
| ELISA (Enzyme-linked Immunosorbent Assay) | 1:10000-30000 |

Product Information

| | |
|-----------------------|--|
| Conjugate | Unconjugate |
| Specificity | Phospho-Catenin δ -1 (Tyr228) Rabbit Polyclonal Antibody detects endogenous levels of catenin δ -1 protein only when phosphorylated at Tyr228. |
| Purification | Affinity purification |
| Concentration | 1mg/ml |
| Format | Liquid |
| Formulation | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol |
| Shipping | Gel Pack |
| Storage | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks |
| UniProt ID | O60716 |
| Entrez-Gene Id | 1500 |

Product Description

This gene encodes a member of the Armadillo protein family, which function in adhesion between cells and signal transduction. Multiple translation initiation codons and alternative splicing result in many different isoforms being translated. Not all of the full-length natures of the described transcript variants have been determined. Read-through transcription also exists between this gene and the neighboring upstream thioredoxin-related transmembrane protein 2 (TMX2) gene.

For Reserch Use Only. Not For Use In Diagnostic Procedures