

**Product Datasheet** 

Order: order@ebiocell.com

TEL: (540)808-3925

Supprt: tech@ebiocell.com
Web: www.ebiocell.com

## Phospho-c-Cbl (Tyr774) Rabbit Polyclonal Antibody

Catalog #: EAB10673

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF/ICC, ELISA	100	Human, Mouse, Rat

## **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting) 1:500-2000
IHC-P(Immunohistochemistry-Paraffin) 1:50-300
IF/ICC(Immunofluorescence/Immunocytochemistry) 1:50-300
ELISA(Enzyme-linked Immunosorbent Assay) 1:5000-20000

## Product Information

Conjugate Unconjugate

Specificity

Phospho-c-Cbl (Tyr774) Rabbit Polyclonal Antibody detects endogenous levels of c-Cbl protein

only when phosphorylated at Tyr774.

**Purification** Affinity purification

Concentration1mg/mlFormatLiquid

Formulation In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol

Shipping Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks

 UniProt ID
 P22681

 Entrez-Gene Id
 867

## **Product Description**

Cbl is a proto-oncogene that encodes a RING finger E3 ubiquitin ligase. The encoded protein is one of the enzymes required for targeting substrates for degradation by the proteasome. This protein mediates the transfer of ubiquitin from ubiquitin conjugating enzymes (E2) to specific substrates. This protein also contains an N-terminal phosphotyrosine binding domain that allows it to interact with numerous tyrosine-phosphorylated substrates and target them for proteasome degradation. As such it functions as a negative regulator of many signal transduction pathways. This gene has been found to be mutated or translocated in many cancers including acute myeloid leukaemia, and expansion of CGG repeats in the 5' UTR has been associated with Jacobsen syndrome. Mutations in this gene are also the cause of Noonan syndrome-like disorder.