

Cortactin Rabbit Polyclonal Antibody

Catalog #: EAB10697

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF/ICC, ELISA	62	Human, Mouse, Rat, Monkey

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

	WB(Western Blotting)	1:500-2000
	IHC-P(Immunohistochemistry-Paraffin)	1:50-300
ELISA(Enzyme-linked Immunosorbent Assay) 1:5000-20000	IF/ICC(Immunofluorescence/Immunocytochemistry)	1:50-300
	ELISA(Enzyme-linked Immunosorbent Assay)	1:5000-20000

Product Information

Conjugate	Unconjugate
Specificity	Cortactin Rabbit Polyclonal Antibody detects endogenous levels of Cortactin protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	<u>Q14247</u>
Entrez-Gene Id	<u>13043</u>

Product Description

This gene is overexpressed in breast cancer and squamous cell carcinomas of the head and neck. The encoded protein is localized in the cytoplasm and in areas of the cell-substratum contacts. This gene has two roles: (1) regulating the interactions between components of adherens-type junctions and (2) organizing the cytoskeleton and cell adhesion structures of epithelia and carcinoma cells. During apoptosis, the encoded protein is degraded in a caspase-dependent manner. The aberrant regulation of this gene contributes to tumor cell invasion and metastasis. Three splice variants that encode different isoforms have been identified for this gene.

For Reserch Use Only. Not For Use In Diagnostic Procedures

EbioCell Lifescineces, Inc.

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA