

## Phospho-FoxM1 (Thr600) Rabbit Polyclonal Antibody

# Catalog #: EAB10762

| Host/Isotype | Clonality  | Applications | MW (kDa) | Reactivity        |
|--------------|------------|--------------|----------|-------------------|
| Rabbit IgG   | Polyclonal | WB           | 84       | Human, Mouse, Rat |

### **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)

1:500-2000

#### **Product Information**

| Conjugate      | Unconjugate   |  |  |
|----------------|---|--|--|
| Specificity    | Phospho-FoxM1 (Thr600) Rabbit Polyclonal Antibody detects endogenous levels of FoxM1 protein only when phosphorylated at Thr600.          |  |  |
| Purification   | Affinity purification   |  |  |
| Concentration  | 1mg/ml  |  |  |
| Format         | Liquid  |  |  |
| Formulation    | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol  |  |  |
| Shipping       | Gel Pack  |  |  |
| Storage        | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles.<br>Aliquots may be stored at +4°C for 1-2 weeks |  |  |
| UniProt ID     | <u>Q08050</u>   |  |  |
| Entrez-Gene Id | <u>2305</u>   |  |  |
|                |   |  |  |

### **Product Description**

FoxM1, also known as FKHL16, MPP2 or Trident, is primarily expressed in proliferating cells. The gene encoding human FOXM1 maps to chromosome 12p13. The transcription element that restricts FOXM1 expression to proliferating cells is located 300 bp upstream of the start codon. FOXM1 is most abundant in thymus, testis, small intestine and colon. Alternative splicing generates FOXM1A and FOXM1B isoforms that contain PEST regions involved in rapid protein degradation. A decrease in FOXM1 expression is associated with age-related defects in cellular proliferation. Conversely, an increase in FOXM1B expression in the livers of older transgenic mice restore hepatocyte DNA replication rates to the higher rate present in young livers. FOXM1B activates the transcription of cyclin B1, cyclin D1 and Cdc25B.

For Reserch Use Only. Not For Use In Diagnostic Procedures

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA