

#### **Product Datasheet**

Order: order@ebiocell.com

Supprt:

Web:

TEL: (540)808-3925 tech@ebiocell.com

www.ebiocell.com

# E-Cadherin Rabbit Polyclonal Antibody

Catalog #: EAB10960

| Host/Isotype | Clonality  | Applications | MW (kDa) | Reactivity |
|--------------|------------|--------------|----------|------------|
| Rabbit IgG   | Polyclonal | WB, IHC-P    | 97       | Human      |

## **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**WB**(Western Blotting) 1:1000-5000 **IHC-P**(Immunohistochemistry-Paraffin) 1:100-500

### **Product Information**

**Conjugate** Unconjugate

**Specificity** E-Cadherin Rabbit Polyclonal Antibody detects endogenous levels of E-Cadherin protein.

**Purification** Affinity purification

Concentration1mg/mlFormatLiquid

Formulation In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol

Shipping Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks

 UniProt ID
 P12830

 Entrez-Gene Id
 999

## **Product Description**

This gene encodes a classical cadherin of the cadherin superfamily. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate the mature glycoprotein. This calcium-dependent cell-cell adhesion protein is comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Mutations in this gene are correlated with gastric, breast, colorectal, thyroid and ovarian cancer. Loss of function of this gene is thought to contribute to cancer progression by increasing proliferation, invasion, and/or metastasis. The ectodomain of this protein mediates bacterial adhesion to mammalian cells and the cytoplasmic domain is required for internalization. This gene is present in a gene cluster with other members of the cadherin family on chromosome 16.