

Cdc25A Rabbit Polyclonal Antibody

Catalog #: EAB11576

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF/ICC, ELISA	59	Human, Mouse, Rat, Monkey

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

IHC-P(Immunohistochemistry-Paraffin)1:50-300IF/ICC(Immunofluorescence/Immunocytochemistry)1:50-300ELISA(Enzyme-linked Immunosorbent Assay)1:5000-20000	WB(Western Blotting)	1:500-2000
	IHC-P(Immunohistochemistry-Paraffin)	1:50-300
ELISA(Enzyme-linked Immunosorbent Assay) 1:5000-20000	IF/ICC(Immunofluorescence/Immunocytochemistry)	1:50-300
	ELISA(Enzyme-linked Immunosorbent Assay)	1:5000-20000

Product Information

Conjugate	Unconjugate
Specificity	Cdc25A Rabbit Polyclonal Antibody detects endogenous levels of Cdc25A protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	<u>P30304</u>
Entrez-Gene Id	<u>993</u>

Product Description

CDC25A is a member of the CDC25 family of phosphatases. CDC25A is required for progression from G1 to the S phase of the cell cycle. It activates the cyclin-dependent kinase CDC2 by removing two phosphate groups. CDC25A is specifically degraded in response to DNA damage, which prevents cells with chromosomal abnormalities from progressing through cell division. CDC25A is an oncogene, although its exact role in oncogenesis has not been demonstrated. Two transcript variants encoding different isoforms have been found for this gene.

For Reserch Use Only. Not For Use In Diagnostic Procedures

EbioCell Lifescineces, Inc.

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA