

Product Datasheet

Order: order@ebiocell.com

TEL: (540)808-3925

Supprt: tech@ebiocell.com
Web: www.ebiocell.com

DBC1 Rabbit Polyclonal Antibody

Catalog #: EAB13276

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF, ELISA	103	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)1:500-2000IHC-P(Immunohistochemistry-Paraffin)1:50-300IF(Immunofluorescence)1:50-300ELISA(Enzyme-linked Immunosorbent Assay)1:5000-20000

Product Information

Conjugate Unconjugate

Specificity DBC1 Rabbit Polyclonal Antibody detects endogenous levels of DBC1 protein.

Purification Affinity purification

Concentration1mg/mlFormatLiquid

Formulation In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol.

Shipping Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks.

 UniProt ID
 Q8N163

 Entrez-Gene ID
 57805

Product Description

DBC-1 (deleted in breast cancer gene 1 protein), also known as p30 DBC protein, is one of the genes located within the region of chromosome 8 (8p21-8p23) that is homozygously deleted in some breast cancers. DBC-1 contains a nuclear localization signal, an N-terminal leucine zipper, an EF hand and a C-terminal coiled-coil region. DBC-1 is closely related to DIS but lacks the SAP domain. During death signaling mediated by TNF α , endogenous DBC-1 undergoes caspase-dependent processing to generate DBC-1 p120 and p66, both of which include the C-terminus of the protein. Both DBC-1 p120 and p66 relocate to the cytoplasm. Overexpression of the DBC-1 p120 form results in mitochondrial clustering and matrix condensation and increases the sensitivity of cells to TNF α -mediated apoptosis. In addition, DBC-1 directly interacts with unliganded ER α , stabilizing its expression and therefore collaborating to suppress apoptosis and promote hormone-independent cell growth.