

Phospho-SHIP2 (Tyr1135) Rabbit Polyclonal Antibody

Catalog #: EAB13348

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF/ICC, ELISA	139	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB (Western Blotting)	1:500-2000
IHC-P (Immunohistochemistry-Paraffin)	1:50-300
IF/ICC (Immunofluorescence/Immunocytochemistry)	1:50-300
ELISA (Enzyme-linked Immunosorbent Assay)	1:5000-20000

Product Information

Conjugate	Unconjugate
Specificity	Phospho-SHIP2 (Tyr1135) Rabbit Polyclonal Antibody detects endogenous levels of SHIP2 when phosphorylated at Tyr1135.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	O15357
Entrez-Gene Id	3636

Product Description

SHIP-2, a homolog of SHIP, is expressed in both hemopoietic and non-hemopoietic cells. In addition to T and B cells, spleen, thymus and lung are shown to coexpress SHIP and SHIP-2. SHIP is also expressed in fibroblasts, heart, skeletal muscle and different brain areas and its expression is enhanced in TSH and EGF-stimulated cells. Like SHIP, SHIP-2 is tyrosine-phosphorylated and associates with Shc after ligation of the B-cell receptor to FcγRII. SHIP-2 causes cell cycle arrest in G1 phase in glioblastoma cells and plays a negative role in regulating the PI 3-kinase-PI 3-kinase B pathway. Both SHIP and SHIP-2 mediate FcγRIIB signaling, including inhibition of proliferation.