

Bcl-w Rabbit Polyclonal Antibody

Catalog #: EAB13364

| Host/Isotype | Clonality | Applications | MW (kDa) | Reactivity |
|--------------|------------|--------------------------|----------|-------------------|
| Rabbit IgG | Polyclonal | WB, IHC-P, IF/ICC, ELISA | 25 | Human, Mouse, Rat |

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| IHC-P(Immunohistochemistry-Paraffin)1:50-300IF/ICC(Immunofluorescence/Immunocytochemistry)1:50-300ELISA(Enzyme-linked Immunosorbent Assay)1:5000-20000 | WB(Western Blotting) | 1:500-2000 |
|--|--|--------------|
| | IHC-P(Immunohistochemistry-Paraffin) | 1:50-300 |
| ELISA(Enzyme-linked Immunosorbent Assay) 1:5000-20000 | IF/ICC(Immunofluorescence/Immunocytochemistry) | 1:50-300 |
| | ELISA(Enzyme-linked Immunosorbent Assay) | 1:5000-20000 |

Product Information

| Conjugate | Unconjugate |
|----------------|---|
| Specificity | Bcl-w Rabbit Polyclonal Antibody detects endogenous levels of Bcl-w protein. |
| Purification | Affinity purification |
| Concentration | 1mg/ml |
| Format | Liquid |
| Formulation | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol |
| Shipping | Gel Pack |
| Storage | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks |
| UniProt ID | <u>Q92843</u> |
| Entrez-Gene Id | <u>599</u> |

Product Description

This gene encodes a member of the BCL-2 protein family. The proteins of this family form hetero- or homodimers and act as anti- and proapoptotic regulators. Expression of this gene in cells has been shown to contribute to reduced cell apoptosis under cytotoxic conditions. Studies of the related gene in mice indicated a role in the survival of NGF- and BDNF-dependent neurons. Mutation and knockout studies of the mouse gene demonstrated an essential role in adult spermatogenesis. Alternative splicing results in multiple transcript variants. Read-through transcription also exists between this gene and the neighboring downstream PABPN1 (poly(A) binding protein, nuclear 1) gene.

For Reserch Use Only. Not For Use In Diagnostic Procedures

EbioCell Lifescineces, Inc.

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA