

## Merlin Rabbit Polyclonal Antibody

### Catalog #: EAB13449

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IF	70	Human

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
<b>IF</b> (Immunofluorescence)	1:50-300

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	Merlin Rabbit Polyclonal Antibody detects endogenous levels of Merlin protein.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
<b>UniProt ID</b>	<a href="#">P35240</a>
<b>Entrez-Gene ID</b>	<a href="#">4771</a>

### Product Description

This gene encodes a protein that is similar to some members of the ERM (ezrin, radixin, moesin) family of proteins that are thought to link cytoskeletal components with proteins in the cell membrane. This gene product has been shown to interact with cell-surface proteins, proteins involved in cytoskeletal dynamics and proteins involved in regulating ion transport. This gene is expressed at high levels during embryonic development; in adults, significant expression is found in Schwann cells, meningeal cells, lens and nerve. Mutations in this gene are associated with neurofibromatosis type II which is characterized by nervous system and skin tumors and ocular abnormalities. Two predominant isoforms and a number of minor isoforms are produced by alternatively spliced transcripts.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**