

EAAT3 Rabbit Polyclonal Antibody

Catalog #: EAB13513

| Host/Isotype | Clonality | Applications | MW (kDa) | Reactivity |
|--------------|------------|--------------|----------|-------------------|
| Rabbit IgG | Polyclonal | WB, ELISA | 57 | Human, Mouse, Rat |

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| | |
|--|--------------|
| WB (Western Blotting) | 1:500-2000 |
| ELISA (Enzyme-linked Immunosorbent Assay) | 1:5000-20000 |

Product Information

| | |
|-----------------------|---|
| Conjugate | Unconjugate |
| Specificity | EAAT3 Rabbit Polyclonal Antibody detects endogenous levels of EAAT3 protein. |
| Purification | Affinity purification |
| Concentration | 1mg/ml |
| Format | Liquid |
| Formulation | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol. |
| Shipping | Gel Pack |
| Storage | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks. |
| UniProt ID | P43005 |
| Entrez-Gene ID | 6505 |

Product Description

EAAT3 (also designated Excitatory amino acid transporter 3, SLC1A1) is a member of the high-affinity glutamate transporters that play an essential role in transporting glutamate across plasma membranes. In brain, these transporters are crucial in terminating the postsynaptic action of the neurotransmitter glutamate, and in maintaining extracellular glutamate concentrations below neurotoxic levels. This transporter also transports aspartate, and mutations in this gene are thought to cause dicarboxylic amino aciduria, also known as glutamate-aspartate transport defect.

For Reserch Use Only. Not For Use In Diagnostic Procedures