

## **Annexin A1 Rabbit Polyclonal Antibody**

# Catalog #: EAB13516

| Host/Isotype | Clonality  | Applications      | MW (kDa) | Reactivity        |
|--------------|------------|-------------------|----------|-------------------|
| Rabbit IgG   | Polyclonal | WB, IHC-P, IF/ICC | 38       | Human, Mouse, Rat |

#### **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| WB(Western Blotting)                           | 1:500-2000 |
|--|------------|
| IHC-P(Immunohistochemistry-Paraffin)           | 1:50-300   |
| IF/ICC(Immunofluorescence/Immunocytochemistry) | 1:50-300   |

#### **Product Information**

| Conjugate      | Unconjugate  |
|----------------|--|
| Specificity    | Annexin A1 Rabbit Polyclonal Antibody detects endogenous levels of Annexin A1 protein.   |
| Purification   | Affinity purification  |
| Concentration  | 1mg/ml   |
| Format         | Liquid   |
| Formulation    | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol.  |
| Shipping       | Gel Pack   |
| Storage        | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles.<br>Aliquots may be stored at +4°C for 1-2 weeks. |
| UniProt ID     | <u>P04083</u>  |
| Entrez-Gene ID | <u>301</u>   |

### **Product Description**

The annexin family of calcium-binding proteins is composed of at least ten mammalian genes and is characterized by a conserved core domain which binds phospholipids in a Ca2+-dependent manner and a unique amino terminal region which may confer binding specificity. The interaction between these proteins and biological membranes have led to the hypothesis that they are involved in cellular trafficking processes such as endocytosis, exocytosis and cellular adhesion. Annexin I, alternatively referred to as lipocortin, has been implicated as a mediator of the anti-inflammatory response produced by glucocorticoids and as an inhibitor of cPLA2, a potent mediator of inflammation. Annexin II, also called p36, has been shown to exist as a monomer or a heterotetramer, complexed with the S-100-related protein p11. This complex is termed calpactin I. In the tetrameric form, Annexin II is an efficient substrate of the PKC family and Src pp60.

For Reserch Use Only. Not For Use In Diagnostic Procedures

EbioCell Lifescineces, Inc.

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA