

Cyclin A2 Rabbit Polyclonal Antibody

Catalog #: EAB13799

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF/ICC	49	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB (Western Blotting)	1:500-2000
IHC-P (Immunohistochemistry-Paraffin)	1:50-300
IF/ICC (Immunofluorescence/Immunocytochemistry)	1:50-300

Product Information

Conjugate	Unconjugate
Specificity	Cyclin A2 Rabbit Polyclonal Antibody detects endogenous levels of Cyclin A2 protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	P20248
Entrez-Gene Id	890

Product Description

Cyclin A2 is a member of the cyclin AB subfamily, cyclin family of proteins. Cyclin A2 is ubiquitously expressed, in contrast to CyclinA1 that only appears in late pachytene spermatocytes. It associates with CDK1 during G2 and M phase, and CDK2 in S phase, of the cell cycle. Binding to SCAPER restricts Cyclin A2 to the cytoplasm. In both compartments, it provides substrate specificity to a phosphorylating complex that promotes entry into, and progression through, mitosis.

For Reserch Use Only. Not For Use In Diagnostic Procedures