

## CTCF Rabbit Polyclonal Antibody

### Catalog #: EAB13867

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IP, IHC-P, IF/ICC, ChIP	83	Human, Mouse, Rat, Monkey

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
<b>IP</b> (Immunoprecipitation)	1:20-200
<b>IHC-P</b> (Immunohistochemistry-Paraffin)	1:50-300
<b>IF/ICC</b> (Immunofluorescence/Immunocytochemistry)	1:50-300
<b>ChIP</b> (Chromatin Immunoprecipitation)	1:10-100

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	CTCF Rabbit Polyclonal Antibody detects endogenous levels of CTCF protein.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol.
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks.
<b>UniProt ID</b>	<a href="http://www.uniprot.org/entry/P49711">P49711</a>
<b>Entrez-Gene ID</b>	<a href="http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&amp;db=Gene&amp;list_uids=10664">10664</a>

### Product Description

This gene is a member of the BORIS + CTCF gene family and encodes a transcriptional regulator protein with 11 highly conserved zinc finger (ZF) domains. This nuclear protein is able to use different combinations of the ZF domains to bind different DNA target sequences and proteins. Depending upon the context of the site, the protein can bind a histone acetyltransferase (HAT)-containing complex and function as a transcriptional activator or bind a histone deacetylase (HDAC)-containing complex and function as a transcriptional repressor. If the protein is bound to a transcriptional insulator element, it can block communication between enhancers and upstream promoters, thereby regulating imprinted expression. Mutations in this gene have been associated with invasive breast cancers, prostate cancers, and Wilms' tumors. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**