

## Fumarase/FH Rabbit Polyclonal Antibody

### Catalog #: EAB13881

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IF/ICC	50	Human, Mouse, Rat

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IF/ICC(Immunofluorescence/Immunocytochemistry)	1:50-300

### Product Information

Conjugate	Unconjugate
Specificity	Fumarase/FH Rabbit Polyclonal Antibody detects endogenous levels of Fumarase/FH protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol.
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks.
UniProt ID	<a href="#">P07954</a>
Entrez-Gene ID	<a href="#">2271</a>

### Product Description

The protein encoded by this gene is an enzymatic component of the tricarboxylic acid (TCA) cycle, or Krebs cycle, and catalyzes the formation of L-malate from fumarate. It exists in both a cytosolic form and an N-terminal extended form, differing only in the translation start site used. The N-terminal extended form is targeted to the mitochondrion, where the removal of the extension generates the same form as in the cytoplasm. It is similar to some thermostable class II fumarases and functions as a homotetramer. Mutations in this gene can cause fumarase deficiency and lead to progressive encephalopathy.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**