

## **SOCS1 Rabbit Polyclonal Antibody**

# Catalog #: EAB13995

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF/ICC, ELISA	24	Human, Mouse, Rat

#### **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IHC-P(Immunohistochemistry-Paraffin)	1:50-300
IF/ICC(Immunofluorescence/Immunocytochemistry)	1:50-300
ELISA(Enzyme-linked Immunosorbent Assay)	1:5000-20000

#### **Product Information**

Conjugate	Unconjugate
Specificity	SOCS1 Rabbit Polyclonal Antibody detects endogenous levels of SOCS1 protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol.
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks.
UniProt ID	<u>O15524</u>
Entrez-Gene ID	<u>8651</u>

### **Product Description**

This gene encodes a member of the STAT-induced STAT inhibitor (SSI), also known as suppressor of cytokine signaling (SOCS), family. SSI family members are cytokine-inducible negative regulators of cytokine signaling. The expression of this gene can be induced by a subset of cytokines, including IL2, IL3 erythropoietin (EPO), CSF2/GM-CSF, and interferon (IFN)-gamma. The protein encoded by this gene functions downstream of cytokine receptors, and takes part in a negative feedback loop to attenuate cytokine signaling. Knockout studies in mice suggested the role of this gene as a modulator of IFN-gamma action, which is required for normal postnatal growth and survival.

For Reserch Use Only. Not For Use In Diagnostic Procedures

EbioCell Lifescineces, Inc.

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA