

CDCP1/CD318 Rabbit Polyclonal Antibody

Catalog #: EAB14005

| Host/Isotype | Clonality | Applications | MW (kDa) | Reactivity |
|--------------|------------|--------------|----------|-------------------|
| Rabbit IgG | Polyclonal | WB, IHC-P | 93 | Human, Mouse, Rat |

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| WB(Western Blotting) | 1:500-2000 |
|--------------------------------------|------------|
| IHC-P(Immunohistochemistry-Paraffin) | 1:50-300 |

Product Information

| Conjugate | Unconjugate |
|----------------|---|
| Specificity | CDCP1/CD318 Rabbit Polyclonal Antibody detects endogenous levels of CDCP1/CD318 protein. |
| Purification | Affinity purification |
| Concentration | 1mg/ml |
| Format | Liquid |
| Formulation | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol |
| Shipping | Gel Pack |
| Storage | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks |
| UniProt ID | <u>Q9H5V8</u> |
| Entrez-Gene Id | <u>64866</u> |

Product Description

CDCP1 (CUB-domain-containing protein 1) contains three extracellular CUB domains, a transmembrane domain, and two putative cytoplasmic tyrosine phosphorylation sites. Phosphorylation of the gp140 and p80 proteins is mediated by Src family kinases at various tyrosine residues, including Tyr 734. PTP family members mediate the desphosphorylation of CDCP1. The conversion of gp140 to p80 prolongs the phosphorylation state, which may affect signaling in epithelial wounds. CDCP1 acts as a marker for hematopoetic cells and also exhibits high expression in metastatic colon and breast tumors.

For Reserch Use Only. Not For Use In Diagnostic Procedures

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA