

#### **Product Datasheet**

Order: order@ebiocell.com

TEL: (540)808-3925

tech@ebiocell.com Supprt: Web:

www.ebiocell.com

# **HAO1** Rabbit Polyclonal Antibody

Catalog #: EAB14143

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF/ICC	41	Human, Mouse, Rat

## **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

1:500-2000 **WB**(Western Blotting) IHC-P(Immunohistochemistry-Paraffin) 1:50-300 IF/ICC(Immunofluorescence/Immunocytochemistry) 1:50-300

### **Product Information**

Conjugate Unconjugate

Specificity HAO1 Rabbit Polyclonal Antibody detects endogenous levels of HAO1 protein.

**Purification** Affinity purification

Concentration 1mg/ml **Format** Liquid

**Formulation** In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol.

Gel Pack Shipping

Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Storage

Aliquots may be stored at +4°C for 1-2 weeks.

**UniProt ID** Q9UJM8 **Entrez-Gene ID** 54363

# **Product Description**

This gene is one of three related genes that have 2-hydroxyacid oxidase activity yet differ in encoded protein amino acid sequence, tissue expression and substrate preference. Subcellular location of the encoded protein is the peroxisome. Specifically, this gene is expressed primarily in liver and pancreas and the encoded protein is most active on glycolate, a two-carbon substrate. The protein is also active on 2hydroxy fatty acids. The transcript detected at high levels in pancreas may represent an alternatively spliced form or the use of a multiple near-consensus upstream polyadenylation site.