EbioCell
LIFE SCIENCES

Product Datasheet

Order: order@ebiocell.com
TEL: (540)808-3925
Supprt: tech@ebiocell.com
Web: *www.ebiocell.com*

Phospho-FoxO4 (Ser262) Rabbit Polyclonal Antibody

Catalog #: EAB14402

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IHC-P, IF, ELISA	54	Human, Mouse

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IHC-P(Immunohistochemistry-Paraffin)	1:50-300
IF(Immunofluorescence)	1:50-300
ELISA(Enzyme-linked Immunosorbent Assay)	1:5000-20000

Product Information

Conjugate	Unconjugate
Specificity	Phospho-FoxO4 (Ser262) Rabbit Polyclonal Antibody detects endogenous levels of FoxO4 protein only when phosphorylated at Ser262.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	P98177
Entrez-Gene Id	4303

Product Description

FoxO4, also known as AFX1, is expressed in a wide variety of tissues and, like other FKHR proteins, AFX1 contains a single forkhead domain and serine-proline-rich region, which mediate DNA binding. AFX1-mediated transcriptional activation is regulated by the serine/threonine kinase Akt1, which phosphorylates AFX1 and in turn, sequesters AFX1 in the cytosol, thereby blocking nuclear localization and DNA binding. Genetic mutations in FKHR genes, including the t(2;13) and t(1;3) translocations, are commonly found in alveolar rhabdomyosarcomas. Additionally, the t(x;11) translocation of the AFX1 gene, which involves the fusion of a serine-proline-rich sequence of AFX1 to the carboxy terminus of a truncated MLL, results in acute lymphocytic leukemia.