

## eIF4A1 Mouse Monoclonal Antibody

### Catalog #: EAB14487

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Mouse IgG1	Monoclonal	WB, IHC-P, IF/ICC	48	Human, Mouse, Rat

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:1000-5000
<b>IHC-P</b> (Immunohistochemistry-Paraffin)	1:50-300
<b>IF/ICC</b> (Immunofluorescence/Immunocytochemistry)	1:50-300

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	eIF4A1 Mouse Monoclonal Antibody detects endogenous levels of eIF4A1 protein.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol.
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks.
<b>UniProt ID</b>	<a href="#">P60842</a>
<b>Entrez-Gene ID</b>	<a href="#">1973</a>

### Product Description

The eukaryotic initiation factor-4A (eIF4A) family consists of 3 closely related proteins EIF4A1, EIF4A2, and EIF4A3. These factors are required for the binding of mRNA to 40S ribosomal subunits. Eukaryotic initiation factor (eIF) complex 2 forms a ternary complex with GTP and the initiator Met-tRNA – this process is regulated by guanine nucleotide exchange and phosphorylation and serves as the main regulatory element of the bottleneck of protein expression.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**