

LAT Rabbit Polyclonal Antibody

Catalog #: EAB14562

| Host/Isotype | Clonality | Applications | MW (kDa) | Reactivity |
|--------------|------------|------------------|----------|-------------------|
| Rabbit IgG | Polyclonal | WB, IHC-P, ELISA | 28 | Human, Mouse, Rat |

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| | |
|--|--------------|
| WB (Western Blotting) | 1:500-2000 |
| IHC-P (Immunohistochemistry-Paraffin) | 1:50-300 |
| ELISA (Enzyme-linked Immunosorbent Assay) | 1:5000-20000 |

Product Information

| | |
|-----------------------|---|
| Conjugate | Unconjugate |
| Specificity | LAT Rabbit Polyclonal Antibody detects endogenous levels of LAT protein. |
| Purification | Affinity purification |
| Concentration | 1mg/ml |
| Format | Liquid |
| Formulation | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol. |
| Shipping | Gel Pack |
| Storage | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks. |
| UniProt ID | O43561 |
| Entrez-Gene ID | 27040 |

Product Description

The protein encoded by this gene is phosphorylated by ZAP-70/Syk protein tyrosine kinases following activation of the T-cell antigen receptor (TCR) signal transduction pathway. This transmembrane protein localizes to lipid rafts and acts as a docking site for SH2 domain-containing proteins. Upon phosphorylation, this protein recruits multiple adaptor proteins and downstream signaling molecules into multimolecular signaling complexes located near the site of TCR engagement. Alternative splicing results in multiple transcript variants encoding different isoforms.

For Reserch Use Only. Not For Use In Diagnostic Procedures