

## **XRCC4 Mouse Monoclonal Antibody**

# Catalog #: EAB14645

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Mouse IgG1	Monoclonal	WB, IP, IHC-P, IF/ICC	38	Human, Mouse, Rat

#### **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

IP(Immunoprecipitation)1:20-200IHC-P(Immunohistochemistry-Paraffin)1:100-500IF/ICC(Immunofluorescence/Immunocytochemistry)1:50-300	WB(Western Blotting)	1:1000-5000
	IP(Immunoprecipitation)	1:20-200
IF/ICC(Immunofluorescence/Immunocytochemistry) 1:50-300	IHC-P(Immunohistochemistry-Paraffin)	1:100-500
	IF/ICC(Immunofluorescence/Immunocytochemistry)	1:50-300

#### **Product Information**

Conjugate	Unconjugate
Specificity	XRCC4 Mouse Monoclonal Antibody detects endogenous levels of XRCC4 protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol.
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks.
UniProt ID	<u>Q13426</u>
Entrez-Gene ID	<u>7518</u>

### **Product Description**

The protein encoded by this gene functions together with DNA ligase IV and the DNA-dependent protein kinase in the repair of DNA double-strand breaks. This protein plays a role in both non-homologous end joining and the completion of V(D)J recombination. Mutations in this gene can cause short stature, microcephaly, and endocrine dysfunction (SSMED). Alternate transcript variants such as NM\_022406 are unlikely to be expressed in some individuals due to a polymorphism (rs1805377) in the last splice acceptor site.

For Reserch Use Only. Not For Use In Diagnostic Procedures

EbioCell Lifescineces, Inc.

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA