

Ras Rabbit Monoclonal Antibody

Catalog #: EAB21365

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IP, FC	21	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IP(Immunoprecipitation)	1:10-100
FC(Flow Cytometry)	1:10-100

Product Information

Conjugate	Unconjugate	
Specificity	Ras Rabbit Monoclonal Antibody detects endogenous levels of total K-Ras, H-Ras, and N-Ras proteins.	
Purification	Affinity purification	
Concentration	1mg/ml	
Format	Liquid	
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol	
Shipping	Gel Pack	
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks	
UniProt ID	<u>P01111, P01112, P01116</u>	
Entrez-Gene Id	<u>4893, 3265, 3845</u>	

Product Description

Ras superfamily is a protein superfamily of small GTPases, which are all related, to a degree, to the Ras protein subfamily (the key human members of which are KRAS, NRAS, and HRAS). Receptor tyrosine kinases and G protein-coupled receptors activate Ras, which then stimulates the Raf-MEK-MAPK pathway. GTPase-activating proteins (GAP) normally facilitate the inactivation of Ras. However, research studies have shown that in 30% of human tumors, point mutations in Ras prevent the GAP-mediated inhibition of this pathway. The most common oncogenic Ras mutation found in tumors is Gly12 to Asp12 (G12D), which prevents Ras inactivation, possibly by increasing the overall rigidity of the protein. This antibody is predicted to react with H-Ras, N-Ras and K-Ras.

For Reserch Use Only. Not For Use In Diagnostic Procedures

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA