

Histone H2A.X Mouse Monoclonal Antibody

Catalog #: EAB21375

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Mouse IgG1	Monoclonal	WB,IHC, IF/ICC, FC	15	Human

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IHC(Immunohistochemistry)	1:100-500
IF/ICC(Immunofluorescence/Immunocytochemistry)	1:100-500
FC(Flow Cytometry)	1:50-200

Product Information

Conjugate	Unconjugate
Specificity	Histone H2A.X Mouse Monoclonal Antibody detects endogenous levels of Histone H2A.X protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	P16104
Entrez-Gene Id	3014

Product Description

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif.

For Research Use Only. Not For Use In Diagnostic Procedures