

Smad3 Rabbit Monoclonal Antibody

Catalog #: EAB21462

| Host/Isotype | Clonality | Applications | MW (kDa) | Reactivity |
|--------------|------------|-----------------------|----------|-------------------|
| Rabbit IgG | Monoclonal | WB, IHC-P, IF/ICC, FC | 48 | Human, Mouse, Rat |

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| | |
|--|------------|
| WB(Western Blotting) | 1:500-2000 |
| IHC-P(Immunohistochemistry-Paraffin) | 1:50-200 |
| IF/ICC(Immunofluorescence/Immunocytochemistry) | 1:50-200 |
| FC(Flow Cytometry) | 1:10-100 |

Product Information

| | |
|----------------|--|
| Conjugate | Unconjugate |
| Specificity | Smad3 Rabbit Monoclonal Antibody detects endogenous levels of Smad3 protein. |
| Purification | Affinity purification |
| Concentration | 1mg/ml |
| Format | Liquid |
| Formulation | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol |
| Shipping | Gel Pack |
| Storage | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks |
| UniProt ID | P84022 |
| Entrez-Gene Id | 4088 |

Product Description

The SMAD family of proteins are a group of intracellular signal transducer proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. The SMAD3 protein functions in the transforming growth factor-beta signaling pathway, and transmits signals from the cell surface to the nucleus, regulating gene activity and cell proliferation. It also functions as a tumor suppressor. Mutations in this gene are associated with aneurysms-osteoarthritis syndrome and Loeys-Dietz Syndrome 3.

For Reserch Use Only. Not For Use In Diagnostic Procedures