

### **Product Datasheet**

Order: order@ebiocell.com

TEL: (540)808-3925

Supprt: tech@ebiocell.com

Web: www.ebiocell.com

# **Lactate Dehydrogenase Rabbit Monoclonal Antibody**

Catalog #: EAB21514

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IP, IHC-P, IF/ICC, FC	37	Human, Mouse, Rat

## **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IP(Immunoprecipitation)	1:10-100
IHC-P(Immunohistochemistry-Paraffin)	1:50-200
IF/ICC(Immunofluorescence/Immunocytochemistry)	1:50-200
FC(Flow Cytometry)	1:10-100

### **Product Information**

**Conjugate** Unconjugate

Specificity

Lactate Dehydrogenase Rabbit Monoclonal Antibody detects endogenous levels of Lactate

Dehydrogenase protein.

**Purification** Affinity purification

Concentration1mg/mlFormatLiquid

Formulation In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol

Shipping Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks

 UniProt ID
 P00338, P07195

**Entrez-Gene ID** 3939, 3945

#### **Product Description**

The protein encoded by this gene catalyzes the conversion of L-lactate and NAD to pyruvate and NADH in the final step of anaerobic glycolysis. The protein is found predominantly in muscle tissue and belongs to the lactate dehydrogenase family. Mutations in this gene have been linked to exertional myoglobinuria. Multiple transcript variants encoding different isoforms have been found for this gene. The human genome contains several non-transcribed pseudogenes of this gene.