

## Aurora B Rabbit Monoclonal Antibody

### Catalog #: EAB21604

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IP, IHC-P, IF/ICC	39	Human

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
<b>IP</b> (Immunoprecipitation)	1:10-100
<b>IHC-P</b> (Immunohistochemistry-Paraffin)	1:50-200
<b>IF/ICC</b> (Immunofluorescence/Immunocytochemistry)	1:50-200

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	Aurora B Rabbit Monoclonal Antibody detects endogenous levels of Aurora B protein.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
<b>UniProt ID</b>	<a href="#">Q96GD4</a>
<b>Entrez-Gene Id</b>	<a href="#">9212</a>

### Product Description

Aurora kinases belong to a highly conserved family of mitotic serine/threonine kinases with three members identified among mammals: Aurora A, B, and C. Studies on the temporal expression pattern and subcellular localization of Aurora kinases in mitotic cells suggest an association with mitotic structure. Aurora kinase functional influences span from G2 phase to cytokinesis and may be involved in key cell cycle events such as centrosome duplication, chromosome bi-orientation and segregation, cleavage furrow positioning, and ingression.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**