

αE-Catenin Rabbit Monoclonal Antibody

Catalog #: EAB21621

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IP, IHC-P, IF/ICC, FC	100	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB (Western Blotting)	1:500-2000
IP (Immunoprecipitation)	1:10-100
IHC-P (Immunohistochemistry-Paraffin)	1:50-200
IF/ICC (Immunofluorescence/Immunocytochemistry)	1:50-200
FC (Flow Cytometry)	1:10-100

Product Information

Conjugate	Unconjugate
Specificity	αE-Catenin Rabbit Monoclonal Antibody detects endogenous levels of αE-Catenin protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	P35221
Entrez-Gene Id	1495

Product Description

This gene encodes a member of the catenin family of proteins that play an important role in cell adhesion process by connecting cadherins located on the plasma membrane to the actin filaments inside the cell. The encoded mechanosensing protein contains three vinculin homology domains and undergoes conformational changes in response to cytoskeletal tension, resulting in the reconfiguration of cadherin-actin filament connections. Certain mutations in this gene cause butterfly-shaped pigment dystrophy.

For Reserch Use Only. Not For Use In Diagnostic Procedures