

## Cox1 Rabbit Monoclonal Antibody

### Catalog #: EAB21675

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IP, IHC-P, IF/ICC, FC	69	Human, Mouse, Rat

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
<b>IP</b> (Immunoprecipitation)	1:10-100
<b>IHC-P</b> (Immunohistochemistry-Paraffin)	1:50-200
<b>IF/ICC</b> (Immunofluorescence/Immunocytochemistry)	1:50-200
<b>FC</b> (Flow Cytometry)	1:10-100

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	Cox1 Rabbit Monoclonal Antibody detects endogenous levels of Cox1 protein.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
<b>UniProt ID</b>	<a href="#">P23219</a>
<b>Entrez-Gene ID</b>	<a href="#">5742</a>

### Product Description

Cyclooxygenase-1 (COX-1), also known as prostaglandin G/H synthase 1, prostaglandin-endoperoxide synthase 1 or prostaglandin H2 synthase 1, is an enzyme that in humans is encoded by the PTGS1 gene. This is one of two genes encoding similar enzymes that catalyze the conversion of arachinodate to prostaglandin. The encoded protein regulates angiogenesis in endothelial cells, and is inhibited by nonsteroidal anti-inflammatory drugs such as aspirin. The protein may promote cell proliferation during tumor progression. Alternative splicing results in multiple transcript variants.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**