

Phospho-GSK-3 β (Ser 9) Rabbit Monoclonal Antibody

Catalog #: EAB21677

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IHC-P, IF/ICC	47	Human, Mouse

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB (Western Blotting)	1:500-2000
IHC-P (Immunohistochemistry-Paraffin)	1:50-200
IF/ICC (Immunofluorescence/Immunocytochemistry)	1:50-200

Product Information

Conjugate	Unconjugate
Specificity	Phospho-GSK-3 β (Ser 9) Rabbit Monoclonal Antibody detects endogenous levels of GSK-3 β only when phosphorylated at Ser9.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	P49841
Entrez-Gene Id	2932

Product Description

Glycogen synthase kinase 3, or GSK-3, is a serine/threonine, proline-directed kinase involved in a diverse array of signaling pathways, including glycogen synthesis and cellular adhesion, and has been implicated in Alzheimer's disease. Two forms of GSK-3, designated GSK-3 α and GSK-3 β , have been identified and differ in their subcellular localization. Tau, a microtubule-binding protein which serves to stabilize microtubules in growing axons, is found to be hyper-phosphorylated in paired helical filaments (PHF), the major fibrous component of neurofibrillary lesions associated with Alzheimer's disease. Hyperphosphorylation of Tau is thought to be the critical event leading to the assembly of PHF. Six Tau protein isoforms have been identified, all of which are phosphorylated by GSK-3. This presents the possibility that miscues in GSK-3 signaling contribute to the onset of Alzheimer's disease.

For Reserch Use Only. Not For Use In Diagnostic Procedures