

## c-Raf Mouse Monoclonal Antibody

### Catalog #: EAB21699

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Mouse IgG1	Monoclonal	WB, IHC-P, FC, ELISA	73	Human

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IHC-P(Immunohistochemistry-Paraffin)	1:100-500
FC(Flow Cytometry)	1:50-200
ELISA(Enzyme-linked Immunosorbent Assay)	1:5000-20000

### Product Information

Conjugate	Unconjugate
Specificity	c-Raf Mouse Monoclonal Antibody detects endogenous levels of c-Raf protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	<a href="#">P04049</a>
Entrez-Gene Id	<a href="#">5894</a>

### Product Description

This gene is the cellular homolog of viral raf gene (v-raf). The encoded protein is a MAP kinase kinase kinase (MAP3K), which functions downstream of the Ras family of membrane associated GTPases to which it binds directly. Once activated, the cellular RAF1 protein can phosphorylate to activate the dual specificity protein kinases MEK1 and MEK2, which in turn phosphorylate to activate the serine/threonine specific protein kinases, ERK1 and ERK2. Activated ERKs are pleiotropic effectors of cell physiology and play an important role in the control of gene expression involved in the cell division cycle, apoptosis, cell differentiation and cell migration. Mutations in this gene are associated with Noonan syndrome 5 and LEOPARD syndrome 2.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**