

#### **Product Datasheet**

Order: order@ebiocell.com

TEL: (540)808-3925 tech@ebiocell.com

Supprt: tech@ebiocell.com
Web: www.ebiocell.com

# Phospho-c-Raf (Ser43) Rabbit Monoclonal Antibody

Catalog #: EAB21824

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB	73	Human, Mouse, Rat

## **Applications Dilutions**

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

**WB**(Western Blotting) 1:500-2000

### **Product Information**

Conjugate Unconjugate

Specificity

Phospho-c-Raf (Ser43) Rabbit Monoclonal Antibody detects endogenous levels of c-Raf protein

only when phosphorylated at Ser43.

**Purification** Affinity purification

Concentration1mg/mlFormatLiquid

Formulation In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol

**Shipping** Gel Pack

Storage Storag

Aliquots may be stored at +4°C for 1-2 weeks

 UniProt ID
 P04049

 Entrez-Gene Id
 5894

## **Product Description**

This gene is the cellular homolog of viral raf gene (v-raf). The encoded protein is a MAP kinase kinase kinase (MAP3K), which functions downstream of the Ras family of membrane associated GTPases to which it binds directly. Once activated, the cellular RAF1 protein can phosphorylate to activate the dual specificity protein kinases MEK1 and MEK2, which in turn phosphorylate to activate the serine/threonine specific protein kinases, ERK1 and ERK2. Activated ERKs are pleiotropic effectors of cell physiology and play an important role in the control of gene expression involved in the cell division cycle, apoptosis, cell differentiation and cell migration. Mutations in this gene are associated with Noonan syndrome 5 and LEOPARD syndrome 2.