

ABCG2 Rabbit Monoclonal Antibody

Catalog #: EAB21917

| Host/Isotype | Clonality | Applications | MW (kDa) | Reactivity |
|--------------|------------|--------------|----------|------------|
| Rabbit IgG | Monoclonal | WB | 72 | Human |

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting) 1:500-2000

Product Information

| | |
|----------------|---|
| Conjugate | Unconjugate |
| Specificity | ABCG2 Rabbit Monoclonal Antibody detects endogenous levels of ABCG2 protein. |
| Purification | Affinity purification |
| Concentration | 1mg/ml |
| Format | Liquid |
| Formulation | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol |
| Shipping | Gel Pack |
| Storage | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks |
| UniProt ID | Q9UNQ0 |
| Entrez-Gene ID | 9429 |

Product Description

The membrane-associated protein encoded by this gene is included in the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. Alternatively referred to as a breast cancer resistance protein, this protein functions as a xenobiotic transporter which may play a major role in multi-drug resistance. It likely serves as a cellular defense mechanism in response to mitoxantrone and anthracycline exposure. Significant expression of this protein has been observed in the placenta, which may suggest a potential role for this molecule in placenta tissue. Multiple transcript variants encoding different isoforms have been found for this gene.

For Reserch Use Only. Not For Use In Diagnostic Procedures