

## ATP-citrate lyase Mouse Monoclonal Antibody

### Catalog #: EAB21925

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Mouse IgG1	Monoclonal	WB, IHC-P, IF/ICC, FC	120	Human, Mouse, Rat, Monkey

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
<b>IHC-P</b> (Immunohistochemistry-Paraffin)	1:100-500
<b>IF/ICC</b> (Immunofluorescence/Immunocytochemistry)	1:50-300
<b>FC</b> (Flow Cytometry)	1:50-200

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	ATP-citrate lyase Mouse Monoclonal Antibody detects endogenous levels of ATP-citrate lyase protein..
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
<b>UniProt ID</b>	<a href="#">P53396</a>
<b>Entrez-Gene Id</b>	<a href="#">47</a>

### Product Description

ATP-citrate lyase is the primary enzyme responsible for the synthesis of cytosolic acetyl-CoA in many tissues. The enzyme is a tetramer (relative molecular weight approximately 440,000) of apparently identical subunits. It catalyzes the formation of acetyl-CoA and oxaloacetate from citrate and CoA with a concomitant hydrolysis of ATP to ADP and phosphate. The product, acetyl-CoA, serves several important biosynthetic pathways, including lipogenesis and cholesterologenesis. In nervous tissue, ATP citrate-lyase may be involved in the biosynthesis of acetylcholine. Multiple transcript variants encoding distinct isoforms have been identified for this gene.