

## MYST2/HBO1 Mouse Monoclonal Antibody

### Catalog #: EAB21973

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Mouse IgG1	Monoclonal	WB, IHC-P, ELISA	71	Human, Mouse, Monkey

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

<b>WB</b> (Western Blotting)	1:500-2000
<b>IHC-P</b> (Immunohistochemistry-Paraffin)	1:50-300
<b>ELISA</b> (Enzyme-linked Immunosorbent Assay)	1:5000-20000

### Product Information

<b>Conjugate</b>	Unconjugate
<b>Specificity</b>	MYST2/HBO1 Mouse Monoclonal Antibody detects endogenous levels of MYST2/HBO1 protein.
<b>Purification</b>	Affinity purification
<b>Concentration</b>	1mg/ml
<b>Format</b>	Liquid
<b>Formulation</b>	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
<b>Shipping</b>	Gel Pack
<b>Storage</b>	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
<b>UniProt ID</b>	<a href="#">O95251</a>
<b>Entrez-Gene ID</b>	<a href="#">11143</a>

### Product Description

HBO1 is a nuclear protein that is highly expressed in human testis. In addition to binding ORC, HBO1 represses AR (androgen receptor)-mediated transcription by binding AR through its N-terminal transcriptional repression domain. HBO1 may play a role in regulating AR-dependent gene transcription in normal and prostate cancer cells. The protein encoded by this gene is part of the multimeric HBO1 complex, which possesses histone H4-specific acetyltransferase activity. This activity is required for functional replication origins and is involved in transcriptional activation of some genes. In both cases, the acetylation of histone H4 helps unfold chromatin so that the DNA can be accessed and replicated or transcribed.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**