

Lamin A/C Rabbit Monoclonal Antibody

Catalog #: EAB21976

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IHC-P, IF/ICC, FC	74	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IHC-P(Immunohistochemistry-Paraffin)	1:100-500
IF/ICC(Immunofluorescence/Immunocytochemistry)	1:50-200
FC(Flow Cytometry)	1:10-100

Product Information

Conjugate	Unconjugate
Specificity	Lamin A/C Rabbit Monoclonal Antibody detects endogenous levels of Lamin A/C protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	P02545
Entrez-Gene Id	4000

Product Description

The protein encoded by this gene is part of the nuclear lamina, a two-dimensional matrix of proteins located next to the inner nuclear membrane. The lamin family of proteins make up the matrix and are highly conserved in evolution. During mitosis, the lamina matrix is reversibly disassembled as the lamin proteins are phosphorylated. Lamin proteins are thought to be involved in nuclear stability, chromatin structure and gene expression. Vertebrate lamins consist of two types, A and B. Alternative splicing results in multiple transcript variants. Mutations in this gene lead to several diseases: Emery-Dreifuss muscular dystrophy, familial partial lipodystrophy, limb girdle muscular dystrophy, dilated cardiomyopathy, Charcot-Marie-Tooth disease, and Hutchinson-Gilford progeria syndrome.

For Reserch Use Only. Not For Use In Diagnostic Procedures