

TFAM Rabbit Polyclonal Antibody

Catalog #: EAB21978

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Polyclonal	WB, IP, IHC-P, IF/ICC, ChIP	29	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB (Western Blotting)	1:5000-20000
IP (Immunoprecipitation)	1:100-500
IHC-P (Immunohistochemistry-Paraffin)	1:200-1000
IF/ICC (Immunofluorescence/Immunocytochemistry)	1:100-500
ChIP (Chromatin Immunoprecipitation)	1:100-500

Product Information

Conjugate	Unconjugate
Specificity	TFAM Rabbit Polyclonal Antibody detects endogenous levels of TFAM protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	Q00059
Entrez-Gene ID	7019

Product Description

mtTFA (mitochondrial transcription factor A), also known as mtTF1, TFAM, TCF6 (for transcription factor 6-like1), TCF6L2 and tsHMG, is a nuclear-encoded gene product that is imported into the mitochondria. mtTFA is required for many aspects of mitochondrial biogenesis including the replication and transcription of mitochondrial DNA (mtDNA). In mouse, testis-specific mtTFA is missing the mitochondria targeting sequence and is present in the nucleus rather than the mitochondria. This form of mtTFA is located primarily in the nuclei of elongated spermatids and may be involved in the regulation of gene expression of the haploid male genome. During mouse and human spermatogenesis there is a reduction of mtTFA protein levels and a reduction in mtDNA copy number. These features may provide one of the mechanisms by which paternal mtDNA transmission is prevented. mtTFA has been associated with mitochondrial disorder in humans characterized by ocular myopathy, exercise intolerance and muscle wasting.

For Reserch Use Only. Not For Use In Diagnostic Procedures