

IMPDH2 Rabbit Monoclonal Antibody

Catalog #: EAB22084

| Host/Isotype | Clonality | Applications | MW (kDa) | Reactivity |
|--------------|------------|--------------|----------|--------------|
| Rabbit IgG | Monoclonal | WB, IF/ICC | 56 | Human, Mouse |

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| WB(Western Blotting) | 1:500-2000 |
|--|------------|
| IF/ICC(Immunofluorescence/Immunocytochemistry) | 1:50-200 |

Product Information

| Conjugate | Unconjugate |
|----------------|---|
| Specificity | IMPDH2 Rabbit Monoclonal Antibody detects endogenous levels of IMPDH2 protein. |
| Purification | Affinity purification |
| Concentration | 1mg/ml |
| Format | Liquid |
| Formulation | In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol |
| Shipping | Gel Pack |
| Storage | Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks |
| UniProt ID | <u>P12268</u> |
| Entrez-Gene ID | <u>3615</u> |

Product Description

This gene encodes the rate-limiting enzyme in the de novo guanine nucleotide biosynthesis. It is thus involved in maintaining cellular guanine deoxy- and ribonucleotide pools needed for DNA and RNA synthesis. The encoded protein catalyzes the NAD-dependent oxidation of inosine-5'-monophosphate into xanthine-5'-monophosphate, which is then converted into guanosine-5'-monophosphate. This gene is up-regulated in some neoplasms, suggesting it may play a role in malignant transformation.

For Reserch Use Only. Not For Use In Diagnostic Procedures

EbioCell Lifescineces, Inc.

Add: Imperial Business Park 4819 Emperor Boulevard, Suite 408 Durham, NC 27703, USA