

## CD71 Rabbit Monoclonal Antibody

### Catalog #: EAB22106

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IHC-P, FC	85	Human

### Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB(Western Blotting)	1:500-2000
IHC-P(Immunohistochemistry-Paraffin)	1:100-500
FC(Flow Cytometry)	1:10-100

### Product Information

Conjugate	Unconjugate
Specificity	CD71 Rabbit Monoclonal Antibody detects endogenous levels of CD71 protein.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	<a href="#">P02786</a>
Entrez-Gene ID	<a href="#">7037</a>

### Product Description

CD71, also known as the transferrin receptor (TFR), is a type II membrane glycoprotein that exists as a disulfide-linked homodimer of two identical subunits. CD71 binds to two molecules of transferrin and a serum iron-transport protein, and directs the cellular uptake of iron via receptor-mediated endocytosis. CD71 is expressed, typically at high levels, on all proliferating cells, reticulocytes and erythroid precursors. It is not expressed on resting leukocytes, but is upregulated upon activation of lymphocytes, monocytes and macrophages. CD71 is also found on most dividing cells and on brain endothelium.

**For Reserch Use Only. Not For Use In Diagnostic Procedures**