

Non-phospho (Active) β -Catenin (Ser33/37/Thr41) Rabbit Monoclonal Antibody**Catalog #: EAB22161**

Host/Isotype	Clonality	Applications	MW (kDa)	Reactivity
Rabbit IgG	Monoclonal	WB, IHC-P	85	Human, Mouse, Rat

Applications Dilutions

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

WB (Western Blotting)	1:2000-10000
IHC-P (Immunohistochemistry-Paraffin)	1:50-200

Product Information

Conjugate	Unconjugate
Specificity	Non-phospho (Active) β -Catenin (Ser33/37/Thr41) Rabbit Monoclonal Antibody detects endogenous β -catenin protein when residues Ser33, Ser37 and Thr41.
Purification	Affinity purification
Concentration	1mg/ml
Format	Liquid
Formulation	In PBS, pH 7.4, Containing 0.02% sodium azide, 0.5% BSA and 50% Glycerol
Shipping	Gel Pack
Storage	Store at -20°C least 1 year from the date of shipment. Avoid repeated freeze/thaw cycles. Aliquots may be stored at +4°C for 1-2 weeks
UniProt ID	P35222
Entrez-Gene Id	1499

Product Description

The protein encoded by this gene is part of a complex of proteins that constitute adherens junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion between cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes cells to stop dividing once the epithelial sheet is complete. Finally, this protein binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatixoma (PTR), medulloblastoma (MDB), and ovarian cancer. Alternative splicing results in multiple transcript variants.

For Reserch Use Only. Not For Use In Diagnostic Procedures